

---

اسم المقال: مقياس قلق لقاح كوفيد - 19 للمراهقين: مقياس لتقييم القلق المرتبط بلقاحات كوفيد - 19 بين المراهقين في مصر  
اسم الكاتب: نسرین یعقوب محمد خليل، ياسر عبد العظيم عبد الموجود سمك  
رابط ثابت: <https://political-encyclopedia.org/library/10394>  
تاريخ الاسترداد: 2026/04/09 09:29 +03

---

الموسوعة السياسية هي مبادرة أكاديمية غير هادفة للربح، تساعد الباحثين والطلاب على الوصول واستخدام وبناء مجموعات أوسع من المحتوى العلمي العربي في مجال علم السياسة واستخدامها في الأرشيف الرقمي الموثوق به لإغناء المحتوى العربي على الإنترنت. لمزيد من المعلومات حول الموسوعة السياسية - Encyclopedia Political، يرجى التواصل على [info@political-encyclopedia.org](mailto:info@political-encyclopedia.org)

استخدامكم لأرشيف مكتبة الموسوعة السياسية - Encyclopedia Political يعني موافقتك على شروط وأحكام الاستخدام المتاحة على الموقع <https://political-encyclopedia.org/terms-of-use>



جامعة الشارقة  
UNIVERSITY OF SHARJAH

# University of Sharjah Journal of Humanities & Social Sciences

A Refereed Scientific journal



Vol. 22, No. 3  
Rabi' al-Awwal 1447 A.H. / September 2025 A.D.

ISSN : 1996-2339

# The COVID-19 Vaccine Anxiety Scale Teen Version: A Mental health screener for COVID-19 Vaccine Related Anxiety among Teenagers

Nisreen Yachoub Mohammed Khalil<sup>(1)</sup>

Yasser Abdelazim Abdelmawgoud Samak<sup>(2)</sup>

Received on: 08 - 01 - 2025

Accepted on: 18 - 02 - 2025

## Abstract:

**Objective:** The COVID-19 pandemic has had a profound impact on the physical life and psychological health of people worldwide. Not only have adults been affected, but children and adolescents have also suffered. The hope of a return to normal life was raised with the development of vaccines, but misinformation about the effectiveness and safety of the vaccines has created another psychological phenomenon, namely anxiety over the vaccinations themselves. Children are more vulnerable to be affected by such fears and anxiety. To address this problem, a unique scale was developed to measure anxiety among teenagers regarding COVID-19 vaccines, and the scale's psychometrics (validity & reliability) were evaluated.

**Methods:** Scale items were developed and after reaching 30 total items, item pooling and division of the items into five subscales were performed. These subscales measure Emotional, Cognitive, Physiological, Behavioral

---

(1) Faculty of Arts and Humanities - King Abdulaziz University ( Jeddah - Saudi Arabia)  
geopsychologie@gmail.com

(2) Faculty of Arts - Assiut University (Assiut - Egypt)

and Emotion Regulation dimensions. Data was collected from 1296 participants with an age range of 12-18. To check the psychometrics of the scale, Exploratory Factor Analysis, Variance, Rotated Factor Matrix, Construct Reliability and Validity along with Confirmatory Factor Analysis were performed.

Results: For the five dimensions that were developed, the Cronbach alpha reliability coefficients were: Emotional Dimension, 0.797; Cognitive Dimension, 0.857; Physiological Dimension, 0.951; Behavioral Dimension, 0.769; and Emotion Regulation, 0.900.

Conclusions: The developed scale is reliable and valid for measuring vaccination anxiety among teens. Future research should follow up this study with a more expansive population worldwide.

**Keywords:** COVID-19 Vaccine Anxiety Scale, COVID-19 Vaccine Anxiety Scale Teen Version, vaccine, anxiety, reliability, validity, Egypt.

## **Background**

The first wave of COVID-19 originated in the city of Wuhan, China at the end of 2019 and rapidly affected every region of the world. According to the World Health Organization, as of 27 August 2021, there were 215 million reported cases and the number of deaths had reached 4.7 million (WHO, 2021). Many researchers have reported that global contagious disease pandemics made many individuals suffer from fear and anxiety (Chong et al., 2004; Wheaton et al., 2012; Wu et al., 2009; Yip et al., 2010). Among health care workers in Egypt and Saudi Arabia, the disease caused anxiety and stress and even depression and sleep disturbances (Arafa et al., 2021). In some people, pre-existing PTSD, generalized anxiety, health anxiety, stress and death anxiety are predictors of enhanced pandemic anxiety (Wheaton et al., 2012). Nonetheless, some degree of Covid-19 anxiety is common in virtually all people due to reasonable fears and uncertainty. Exacerbating the situation, ignorance and misinformation have added to many individuals' uncertainty and anxiety about the illness (Harper et al., 2020; Banerjee, 2020).

As of this writing, vaccines for COVID-19 are the only way to end this pandemic (Lurie et al., 2020). These vaccines are the most effective way to control, prevent, and build immunity in individuals for the disease. The efficacy of these already developed vaccines for society is effectively dependent on the vaccination rates of individuals. Increasing the rate of those who are vaccinated relies on individuals' willingness, which in turn depends on their attitudes toward and acceptance of the vaccines (Wang et al., 2020). With respect to children and adolescents, getting vaccinated is effectively dependent on parents' acceptance of the vaccine (Lazarus et al., 2021). The usefulness of COVID-19 vaccines against the pandemic will be

compromised if a large percentage of people choose not to get vaccinated (Murray & Piot, 2021). The COVID-19 pandemic led to radical changes in students' lifestyles, as education moved entirely to a virtual environment. This shift significantly impacted the educational process, causing numerous challenges for both students and teachers. (Atiyat & Abu Hamour, 2022).

Recent studies have emphasized the importance of psychometric validation in assessing vaccine-related anxiety. Khusaifan and Samak (2024) developed and validated the COVID-19 Vaccine Anxiety Scale (CVAS) for adults, demonstrating its reliability and validity across different dimensions of vaccine-related anxiety. Their findings highlight the necessity of tailored anxiety scales to address concerns specific to various age groups and demographics, which aligns with the objectives of this study in assessing adolescents' vaccine anxiety. Their research concluded that psychometrically validated scales provide accurate assessments that are crucial for designing mental health interventions related to vaccine hesitancy (Khusaifan & Samak, 2024).

One of the important and effective roles of health authorities must be to create awareness among the public regarding vaccine effectiveness and safety to get them to accept the vaccines and get vaccinated to prevent themselves and their children from getting the disease as well as halting the spread of the disease and the concomitant evolution of new variants. Some governments have assessed the acceptance rate among their citizens. As was reported in China, the acceptance rate is 91.3%, in Brazil, it is 85%, in African countries, mostly in South Africa, it is 79%, with the same rate in South Korea as well. About European countries, in France and Italy the acceptance rate is 62%. Russia has a 55% acceptance rate, while in the United States it is 76% (Neumann-Böhme et al., 2020; Detoc et al.,

2020; Pogue et al., 2020; Szilagyi et al., 2020; Daly & Robinson, 2020). There are still countries with low rates of vaccination acceptance, including some Arab countries, Central African nations, and Asian countries, where people are still rejecting vaccines, with many considering vaccination a serious threat to their health or even lives. In some Arab and African nations, vaccination acceptance is less than 30% (Sallam et al., 2020; Nzaji et al., 2020). When studying health care workers in Pakistan, Malik, Malik, and Ishaq, (2021) found that vaccine acceptance increases with increasing age but among reasons cited for vaccine hesitancy, concerns about vaccine effectiveness and side effects were prevalent. Magadmi and Kamel (2021) reported that among Saudi Arabian people, almost 55.3% had hesitancy and fearful thoughts regarding the Covid vaccinations. In the Egyptian population findings of Saied et al. (2020), it was observed that despite 90% of respondents agreeing that there is a need for vaccination, still more than 46% said they will not accept vaccination at this stage as they are fearful, phobic, and anxious regarding the effectiveness of COVID-19 vaccinations. These adult attitudes surely impact the mindsets of teens and adolescents as at this age, they are influenced by elders' decisions. Also, this age group is more vulnerable to developing anxiety symptoms regarding Covid vaccinations (Barth, 2021). Clearly, many individuals, including children, are facing fear and anxieties regarding vaccinations' consequences to their health. This contributed to improving the mental health of infected individuals and enhancing their ability to cope with the challenges they faced. (Ali, 2024, p. 525). According to Ali and Hassan (2023), The COVID-19 pandemic has significantly impacted all aspects of social and economic life, leading to the suspension of economic activities, the closure of educational institutions, and the imposition of lockdowns on populations in many countries.

Anxiety and fears about the disease and the pandemic can be due to a lack of scientific knowledge. And the same holds true for acceptance of COVID-19 vaccinations as well. This anxiety may expose people to false news about the acceptance and significance of COVID-19 vaccines. Anxiety and fear can destabilize one's immunity and make people more vulnerable to the coronavirus (Bajema et al., 2020, Chan et al., 2020). In similar previous situations (e.g., MERS, SARS), as well as for Covid, measurement scales were developed to measure the consequences of persistent illness, as well as vaccination acceptance, fears, and anxieties. (Pérez-Fuentes et al., 2020; Serrano-Mollar, 2012; Yohannes et al., 2013). In the current situation, it is essential to create valid scales to determine individuals' level of anxiety related to COVID-19 and vaccines, and develop strategies to deal with this situation, such as education on the effectiveness of vaccines against such diseases as well as their safety profile. Sufficient awareness of correct information about COVID-19 and vaccines may help keep people secure, but too much attention can also be psychologically disturbing and harmful (Taylor, 2019). The media also plays a vital role in both regards, disseminating truthful and correct scientific information while potentially triggering higher levels of anxiety and phobias (Kumar & Somani, 2020; Lee, 2020b; Hamza et al., 2020; Bodner et al., 2021)

A recent study of 775 adults in the United States found that people who became anxious about the coronavirus and whose performance was affected in various ways became frustrated, even suicidal, and attempted to cope via religious and alcohol/substance abuse (Lee, 2020a). Therefore, Lee (2020a) suggested that it is essential for healthcare experts to understand their patients' psychological state to help people with coronavirus anxiety and COVID-19 vaccination anxiety. This is particularly important for

individuals in their teenage years because teens can develop anxiety quickly and this may affect them their entire life span (Asmundson & Taylor 2020; Lee,2020b).

Thus, anxiety regarding vaccinations is becoming another global issue in addition to the COVID-19 pandemic itself. It is very important to consider its destructive effects on teens and adolescents. Negative information about vaccinations, trials, errors, failures, and fake news, etc., are making individuals, including those in this age group, more anxious regarding vaccines despite our knowing that this is the best way of survival in this pandemic. Because people are anxious about Covid vaccinations, there is a need for a suitable tool to assess individuals' vaccine anxiety so mental health care providers can design appropriate interventions and programs to reduce their fears. It is necessary to design a scale with good validity and reliability for measuring COVID-19 vaccination-related anxiety among all individuals as well as children and adolescents. This study aimed to determine the psychometric properties of the COVID-19 Vaccination Anxiety Scale Teen Version for individuals in the age range from 12-18 in the Egyptian population.

### **Purpose of the Study**

Researchers have reported on studies of adults regarding COVID-19 vaccination anxiety and acceptance. However, there is a huge gap in the literature regarding younger people. So, the purpose of this research was to address such issues in the teenage population. To fill this gap and provide a reliable and valid screening scale, this work began by developing the COVID-19 Vaccine Anxiety Scale Teen Version (CVAS Teen Version) and then tested it with a sample of teens with an age range from 12-18 in Egypt.

## Methods

For this study, we used a cross-sectional research design for the collection of self-reported response survey data from Egypt-resident teens. Online consent was obtained from the guardians and caregivers of those teen participants to fulfil the ethical consideration. Parents and caregivers were briefly informed of the study's purpose. The work was completed in two phases. In the first phase, item pooling was accomplished, resulting in five different dimensions for this scale. In the second phase, from the targeted sample of Egyptian teens with an age range from 12 to 18, data was collected from 1246 (male=786, female=460) participants. The data collection was performed over five months, from the 1<sup>st</sup> of January 2021 to the 31<sup>st</sup> of May, 2021. Data collection was conducted by having participants respond to questionnaires via Google Forms.

### Item Pooling with Subscale Description

The COVID-19 Vaccination Anxiety Scale Teen Version is a unique scale, so to develop item pooling and division of the items into subdomains, first a systematic review of the literature was conducted. Then, expert opinion was obtained regarding theoretical aspects, literature, analysis, and finalization of scale items. This resulted in the division of items into five subdomains: Emotional Dimension (ED), Cognitive Dimension (CD), Physiological Dimension (PD), Behavioral Dimension (BD), and Emotion Regulation (ER). Responses for each scale item used a Likert scale ranging from 1 to 5 (1=strongly disagree, 2=disagree, 3=neutral, 4=agree and 5=strongly agree).

## Results

The final version of the COVID-19 Vaccination Anxiety Scale Teen Version is shown in the Appendix. Multivariate analyses were performed after specifying the suitable sample size; as per Preacher and MacCallum (2002), it should be not less than 200. In the current study, demographic data was collected via questionnaire. Among 1296 teen participants, 836(65.5%) were male and 460 (34.5%) females, the majority of whom (98.9%) were in the age range of 16-18 (Table1).

**Table 1 Demographical Profile of the Sample, n=1296**

Variable	Category	<i>f</i>	%
Gender	Male	836	% 35.5
	Female	460	% 64.5
Age	Years 13-15	14	1.1%
	Years 16-18	1232	98.9%

## Validating the Scale

This research sought to validate the scale that was used to measure five key dimensions, that is, Emotional Dimension (ED), Cognitive Dimension (CD), Physiological Dimension (PD), Behavioral Dimension (BD), and Emotion Regulation (ER). The latter, Emotion Regulation, was measured by two sub-dimensions, namely Reappraisal (RE) and Suppression (SUP). Being a study that used the scale for a new population, it was imperative, according to Judd, McClelland, and Ryan (2017), to ascertain whether the measurement items explained the dimensions well and this entailed the use of Exploratory Factor Analysis (EFA). To further validate the dimensions, construct validity was conducted using Confirmatory Factor Analysis (CFA).

## Exploratory Factor Analysis

Before the use of EFA, it was imperative to test the adequacy of the used sample. This was done through the Kaiser-Mayer-Olkin (KMO) measure of sampling adequacy (Loehlin& Beaujean, 2017). The results are presented in Table 2.

**Table 2 KMO and Bartlett's Test**

Kaiser-Meyer-Olkin Measure of Sampling Adequacy		953.
Bartlett's Test of Sphericity	Approx. Chi-Square	21784.977
	df	435
	.Sig	000.

The results show that  $KMO = 0.953$  and this was greater than the minimum expected 0.50 (Thompson, 2018), suggesting that the sample size for the sample data used was more than the minimum necessary size. Further, for Bartlett's test of sphericity,  $\chi^2(435) = 62309.337$ ,  $p < 0.05$ , and, again, this meant that the assumption was not violated. Since the focus of the analysis was to extract factors rather than reduce the items, the Principal Axis Factoring (PAF) method was considered instead of Principal Component Analysis (PCA). Further, the constructs were assumed to be marginally correlated or not correlated at all, hence the orthogonal rotation. The importance of each of the items used in the factor analysis was then determined through the computation of the communalities. These indicated the amount of variance in each variable that is accounted for and, according to Little (2013), the minimum variance expected is 0.40. The results are summarized in Table 3.

None of the communalities was less than 0.40, with the minimum

observed being 0.514 for the item: “I experience chest discomfort when I read or listen to COVID-19 vaccine news.” The second lowest item had a communality of 0.520 and this was: “I feel nauseous when I think of getting the COVID-19 vaccine.” The variances are presented in Table 4.

**Table 3 Scale Items and Communalities**

	Initial	Extraction
I find it difficult to relax when I think about taking the COVID-19 vaccine.	.450	.603
The thought of getting a vaccination makes me restless.	.481	.569
Even after information about vaccine trials and success rates, I am not able to control my worry.	.452	.553
I am afraid that the vaccine could cause possible harm to me and my family.	.495	.584
I am scared that the COVID-19 vaccine could cause the disease.	.491	.562
I use to have repetitive negative thoughts about the consequences of COVID	.461	.530
I am fearful of the side effects of the COVID-19 vaccine.	.466	.536
I am scared that the COVID-19 vaccine could cause some sort of deformity or dysfunction in my body.	.456	.525
I experience shortness of breath when I think of taking COVID	.424	.550
I experience chest discomfort when I read or listen to the COVID-19 vaccine news.	.396	.514
I feel nauseous when I think of getting the COVID-19 vaccine.	.392	.520

The COVID-19 Vaccine Anxiety Scale Teen Version: A Mental health screener for COVID-19 Vaccine Related Anxiety among Teenagers (813-852)

I experience loosening of bowels when think of taking the COVID-19 vaccine.	.663	.695
My heart starts to run faster when I think or read about the COVID-19 vaccine.	.658	.691
I feel some sensation in my muscles when I think about getting the COVID-19 vaccine.	.664	.691
I experience pain in my muscles when I think of getting the COVID-19 vaccine.	.680	.713
I experience hot and cold flushes when I think about getting the COVID-19 vaccine.	.651	.678
I feel numb when I think of taking the COVID-19 vaccine.	.647	.676
I frequently check for vaccine validation facts to ensure its effectiveness.	.668	.697
The thought of taking the COVID 19 vaccine causes difficulty in falling asleep and bad dreams.	.622	.646
I would prefer to avoid vaccines in case it's not compulsory	.655	.687
When I want to feel happier, I think about something different	.518	.575
When I want to feel less bad (e.g., sad, angry, worried) I think about something different.	.538	.596
When I'm worried about something; I make myself think about it in a way that helps me feel better.	.509	.566
When I want to feel happier about something, I change the way I'm thinking about it.	.527	.592
I control my feelings about things by changing the way I think about them.	.530	.579

When I want to feel less bad about something I change the way I'm thinking about it.	.537	.600
I keep my feelings to myself.	.511	.606
When I am feeling happy, I am careful not to show it.	.498	.588
I control my feelings by not showing them.	.499	.594
When I'm feeling bad (e.g., sad, angry, worried), I am careful not to show it.	.474	.558
<i>Note.</i> Extraction Method: Principal Axis Factoring.		

**Table 4 Variance Explained by Factors**

Factor	Initial Eigenvalues			Extraction Sums of Squared Loadings			Rotation Sums of Squared Loadings		
	Total	% Of Variance	Cumulative %	Total	% Of Variance	Cumulative %	Total	% Of Variance	Cumulative %
1	9.378	31.261	31.261	9.008	30.028	30.028	6.300	21.001	21.001
2	5.134	17.114	48.375	4.727	15.756	45.784	3.764	12.547	33.548
3	2.254	7.512	55.887	1.852	6.174	51.958	2.991	9.971	43.520
4	1.539	5.129	61.016	1.104	3.680	55.638	2.105	7.016	50.536
5	1.181	3.935	64.951	.737	2.455	58.094	1.504	5.014	55.550
6	1.074	3.580	68.531	.644	2.147	60.240	1.407	4.690	60.240
7	.532	1.774	70.305						
8	.519	1.729	72.034						
9	.500	1.666	73.699						

The COVID-19 Vaccine Anxiety Scale Teen Version: A Mental health screener for COVID-19 Vaccine Related Anxiety among Teenagers (813-852)

10	.493	1.642	75.342						
11	.482	1.606	76.948						
12	.462	1.541	78.489						
13	.458	1.528	80.018						
14	.441	1.470	81.487						
15	.430	1.432	82.920						
16	.424	1.412	84.332						
17	.409	1.364	85.696						
18	.398	1.325	87.021						
19	.393	1.309	88.330						
20	.384	1.279	89.610						
21	.369	1.230	90.840						
22	.363	1.210	92.050						
23	.337	1.122	93.172						
24	.327	1.089	94.261						
25	.312	1.040	95.301						
26	.304	1.013	96.314						
27	.290	.966	97.279						
28	.286	.952	98.232						
29	.274	.915	99.146						
	30								
	.256								
	.854								
	100.000								
<i>Note.</i> Extraction Method: Principal Axis Factoring.									

According to Hoyle (2012) and Hancock and Mueller (2013), the minimum expected eigenvalue is 1.0 and from the results, six factors had eigenvalues that were greater than 1.0. This meant that out of all the items, six factors had been extracted. Factor 1, whose eigenvalue was  $\lambda = 9.378$ , explained the greatest variance of 21.001%, followed by Factor 2 ( $\lambda = 5.134$ ) which explained 12.547% of the variance. Factor 3 ( $\lambda = 2.254$ ) explained 9.971% of the variance, Factor 4 ( $\lambda = 1.539$ ) explained 7.016% of the variance, while Factor 5 ( $\lambda = 1.181$ ) explained 5.014% of the variance. Finally, Factor 6 ( $\lambda = 1.074$ ), explained 4.690% of the variance. The cumulative variance explained was 60.240% and is greater than the minimum criterion of 50%; this confirmed that the extracted factors were valid (Byrne, 2016; Thompson, 2018; Gana and Broc, 2019). Table 5 presents the rotated factor matrix.

The outcomes show that six factors had been extracted. Comparing the items with factor loadings greater than 0.50 (Bandalos & Finney, 2010; Thompson, 2018), it can be seen that for Factor 1, the items measured the *Physiological Dimension*, for Factor 2, they measured *Reappraisal*, for Factor 3, they measured the *Cognitive Dimension*, for the fourth factor, the items measured *Suppression*, for Factor 5, the items measured the *Behavioral Dimension*, and lastly, Factor 6 items measured the *Emotional Dimension*. It should be noted that the EFA result had grouped the dimensions according to the original conceptualized form and this confirms that the scale was good.

### **Construct Reliability**

According to Howitt and Cramer (2017), the reliability of the research constructs ought to be tested before their use in modelling to evaluate

internal consistency. In this regard, it was necessary to ensure that the reliability of the underlying variables used in the research was tested.

**Table 5 Rotated Factor Matrix**

Items	Factor					
	1	2	3	4	5	6
	.232	.041	.279	.095	.104	<b>.671</b>
	.291	.068	.367	.025	.132	<b>.572</b>
	.230	.032	.310	.073	.170	<b>.607</b>
	.240	.038	<b>.702</b>	.026	.139	.108
	.233	.001	<b>.675</b>	.041	.119	.188
	.241	.043	<b>.656</b>	.074	.123	.140
	.239	.051	<b>.657</b>	.058	.130	.156
	.221	.059	<b>.664</b>	.046	.106	.135
	.273	.081	.209	.042	<b>.639</b>	.128
	.276	.119	.166	.001	<b>.619</b>	.111
	.233	.099	.197	-.00	<b>.639</b>	.087
	<b>.797</b>	.006	.173	.059	.138	.086
	<b>.804</b>	.017	.164	.028	.108	.071
	<b>.794</b>	-.01	.191	.032	.115	.099
	<b>.808</b>	-.00	.162	.070	.133	.107
	<b>.770</b>	.002	.206	.015	.178	.106
	<b>.788</b>	-.01	.159	.046	.120	.113
	<b>.791</b>	.000	.185	.032	.148	.121
	<b>.768</b>	.033	.174	.018	.122	.100

	<b>.785</b>	-.02	.212	.059	.074	.131
	.000	<b>.728</b>	.004	.205	.038	.032
	-.019	<b>.737</b>	.069	.212	.041	.012
	-.011	<b>.733</b>	.005	.151	.079	.008
	.009	<b>.754</b>	.013	.150	.030	.002
	-.012	<b>.719</b>	.056	.226	.058	.068
	.010	<b>.750</b>	.045	.173	.069	.009
	.061	.347	.079	<b>.689</b>	.011	.030
	.103	.345	.055	<b>.672</b>	.008	.057
	.064	.327	.040	<b>.690</b>	.012	.073
	.020	.331	.062	<b>.665</b>	.036	.025
<i>Note:</i> Extraction Method: Principal Axis Factoring. Rotation Method: Varimax with Kaiser Normalization. Rotation converged in 6 iterations.						

To achieve this, Cronbach's alpha was computed and, as per Kilic (2008), the minimum acceptable alpha was 0.70. The results are summarized in Table 6.

From the outcome, none of the constructs had a reliability alpha that was less than the minimum criterion of 0.70. The minimum observed alpha was for the construct *Behavioral Dimension* ( $\alpha = 0.744$ ) and thus the results show that the dimensions used met the reliability requirements.

**Table 6 Reliability Testing**

<b>Construct</b>	<b>N of items</b>	<b>Alpha</b>
Emotional Dimension	3	.797
Cognitive Dimension	5	.857
Physiological Dimension	9	.951
Behavioral Dimension	3	.769
Emotion Regulation	10	.900
Reappraisal (RE)	6	.893
Suppression (SUP)	4	.849

### **Test-Retest Reliability**

To confirm test-retest reliability, we used intraclass correlation coefficients (ICCs) and Kappa coefficients (KCs). ICCs are an ideal index to measure the scale’s test-retest reliability with reflecting correlation along with agreement (Portney & Watkins, 2009). To quantify the agreement level, KCs were used on categorical variables for assessing the inter-rated and test-retest reliability for clinical scales (Brenner & Kliedsch, 1996). For all domains of this scale, the average ICC score was 0.901 and average KC was 0.867, which showed considerable test-retest reliability for all five sub-dimensions of the scale (see Table 7).

**Table 7 Test-retest Reliability**

	<b>ED</b>	<b>CD</b>	<b>BD</b>	<b>PD</b>	<b>RE</b>	<b>SUP</b>
ICC	0.891	0.911	0.861	0.902	0.931	0.911
KC	0.856	0.812	0.941	0.821	0.871	0.904

*Note:* ICC=intraclass correlation coefficient, KC= Kappa coefficient

## **Construct Validity**

Having tested the reliability of the constructs, it was also essential to validate whether the constructs were valid. To achieve this, confirmatory factor analysis was also carried out, over and above the exploratory factor analysis, and the measurement model is shown in Figure 1.

The validity of the constructs was tested by measuring both the convergent and discriminant validity. The convergent validity of the constructs was measured using the Average Variance Extracted (AVE), which, according to Hox (2013) and Kline (2016), ought to be at least 0.50. On the other hand, the discriminant validity was tested using the Heterotrait-Monotrait Ratio (HTMT), which, according to Fox et al. (2012), Brown (2015), and Kline (2016), ought to be less than 0.85. The results for both tests are presented in Table 8.

Concerning convergent validity, the minimum AVE was 0.526 for the Reappraisal (RE) construct and because all the AVEs were greater than 0.50, this confirms that the convergent validity assumption was not violated. On the other hand, the highest HTMT score was 0.694, between the Behavioral Dimension (BD) and Suppression (SUP), and because this was less than the maximum tolerable threshold of 0.85, it confirms that the discriminant validity was not violated. Therefore, by and large, the constructs used in the study were reliable and valid.

**Table 8 Convergent and Discriminant Validity**

	CR	AVE	MSV	MaxR(H)	ED	CD	BD	PD	RE	SUP
ED	0.951	0.684	0.309	0.951	0.827					
CD	0.893	0.581	0.422	0.893	0.031	0.762				
BD	0.857	0.545	0.482	0.857	***0.556	***0.120	0.738			
PD	0.849	0.584	0.422	0.849	***0.149	***0.649	***0.191	0.764		
RE	0.769	0.526	0.308	0.770	***0.555	***0.210	***0.539	***0.149	0.725	
SUP	0.796	0.566	0.482	0.797	***0.548	***0.134	***0.694	***0.214	***0.523	0.752

Note. \*\*\*p>.001

*AVE, Average Variance Extracted; CR, Composite Reliability; MaxR(H), maximum reliability; MSV, Maximum Shared Squared Variance.*

**Table 9 Confirmatory Factor Analysis (CFA) Model Fit Summary**

### CMIN

Model	NPAR	CMIN	DF	P	CMIN/DF
Default model	75	437.851	390	.047	1.123
Saturated model	465	.000	0		
Independence model	30	21975.277	435	.000	50.518

### Baseline Comparisons

Model	NFI Delta1	RFI rho1	IFI Delta2	TLI rho2	CFI
Default model	.980	.978	.998	.998	.998
Saturated model	1.000		1.000		1.000
Independence model	.000	.000	.000	.000	.000

### Parsimony-Adjusted Measures


Model	PRATIO	PNFI	PCFI
Default model	.897	.879	.895
Saturated model	.000	.000	.000
Independence model	1.000	.000	.000

### RMSEA

Model	RMSEA	LO 90	HI 90	PCLOSE
Default model	.010	.001	.015	1.000
Independence model	.199	.197	.201	.000

*Table 9 presents the CFA model fit summary, including key goodness-of-fit indices such as CMIN/DF, NFI, RFI, IFI, TLI, CFI, and RMSEA. The results indicate a good model fit, with the default model showing favorable comparative fit indices (CFI = 0.998) a*

**Figure 1 Measurement Model: Confirmatory Factor Analysis**



This structural equation model (SEM) diagram illustrates the relationships between different dimensions of emotional regulation, suppression, and their associated observed indicators. The latent variables include Emotional Dimension (ED), Cognitive Dimension (CD), Behavioral Dimension (BD), Physiological Dimension (PD), Emotion Regulation Reappraisal (RE), and Suppression (SUP). Each latent variable is measured by multiple observed indicators (PD1–PD9, RE1–RE6, CD1–CD5, SUP1–SUP4, BD1–BD3, and ED1–ED3), with standardized factor loadings shown for each. Correlations between latent variables are indicated with curved arrows, highlighting the strength and direction of associations.

## **CFA Model Goodness of Fit**

According to Wang and Wang (2019), it is crucial to evaluate the goodness of fit of the structural equation model. In this respect, several indices were reviewed. These comprised the  $\chi^2/df$  which, according to Fox et al. (2012) and Arbuckle (2016), should be less than 3.0. The second layers were the baseline comparisons, and these included the normed fit index (NFI) as well as the comparative fit index (CFI), both of which ought to be greater than 0.90 (Boomsma, Hoyle, & Panter, 2012; Hoyle, 2012; Arbuckle, 2016). Parsimony adjusted measures, which included the parsimonious normed fit index (PNFI) and the parsimonious comparative fit index (PCFI), were also calculated. These were expected to be greater than 0.50 (Gefen, Rigdon, & Straub, 2011; Wang & Wang, 2019). Lastly, the Root Mean Square Error of Approximation (RMSEA) is based on the non-centrality index and should be less than 0.08. The model fit results from the analysis are presented in Table 9.

From the findings,  $\chi^2/df = 1.123$ , which is  $< 3.0$  and thus was less than the maximum acceptable threshold. For the baseline comparisons,  $NFI = 0.980$ , which is  $> 0.90$ ,  $RFI = 0.978 > 0.90$ ,  $IFI = 0.998 > 0.90$ ,  $TLI = 0.998 > 0.90$  and  $CFI = 0.998 > 0.90$ . Regarding the parsimony adjusted measures,  $PNFI = 0.879 > 0.50$ , while the  $PCFI$  was  $0.895 > 0.50$ . Lastly, the  $RMSEA$  statistic was  $0.010 < 0.08$  and thus was within the desired threshold. The overall conclusion was that the CFA model and the results therefrom were valid.

## **Discussion**

Considering the worldwide COVID-19 pandemic and its psychological effects on individuals, this research aimed to investigate the psychometric properties of the teen version of the COVID-19 Vaccination Anxiety

Scale. The results showed that the CVAS Teen Version with a five-dimensional structure has good psychometric properties.

In this study, to evaluate the convergent validity, the CVAS Teen Version's five sub-dimensions – Emotional Dimension, Cognitive Dimension, Physiological Dimension, Behavioral Dimension, and Emotion Regulation, the latter with two sub-dimensions, Reappraisal and Suppression– were used. The lowest AVE statistic was 0.526 (for the Reappraisal construct) and all the AVEs were greater than 0.50, thus validating that the convergent validity assumption was not violated. The outcomes associated with construct validity also showed that the six-factor structure has a good model fit.

For variance explanation, factor analysis was used. As discussed in the Exploratory Factor Analysis section, the cumulative variance explained was 60.24%, which is greater than the minimum criterion of 50%. This confirmed that the extracted factors were valid (Byrne, 2016; Thompson, 2018; Gana & Broc, 2019).

The CVAS Teen Version demonstrated excellent internal consistency and the Cronbach's alpha coefficients on all five dimensions conformed to the necessary criteria. These results were consistent with Lee's (2020a) main study, which reported Cronbach's alpha of 0.93 for the Coronavirus Anxiety Scale, and were in line with the results of the COVID-19 vaccination acceptance scale (Esteban et al., 2021) which was 0.83. Also, all items had good content validity, indicating that all questions were related to anxiety about COVID-19 vaccination for the age range of 12-18. Construct reliability was also confirmed on all five subdomains of the CVAS Teen Version as it was found that the Emotional Dimension's  $\alpha = 0.797$ , the

Cognitive Dimension's  $\alpha = 0.857$ , the Physiological Dimension's  $\alpha = 0.951$ , the Behavioral Dimension's  $\alpha = 0.769$  and Emotion Regulation's  $\alpha = 0.900$ , while for Emotional Regulation's two sub-dimensions, Reappraisal had  $\alpha = 0.893$  and Suppression had  $\alpha = 0.849$ . Also, test-retest reliability using ICC and KC was acceptable. These findings confirmed that the reliability of the research constructs ought to be tested before their use in modelling to evaluate the internal consistency (Howitt & Cramer, 2017). In this regard, none of the constructs had a reliability alpha that was less than the minimum criterion of 0.70. The results showed that the dimensions used met the reliability requirements.

The outcomes of this research study are consistent with those of Evren et al. (2020). In their research, the Cronbach's alpha for the Coronavirus Anxiety Scale (CAS) was reported to be 0.80. In addition, CAS showed a positive correlation with the Fear of COVID-19 Scale (Ahorsu et al., 2020) and the Corona Obsession Scale (Lee, 2020b). The diagnostic accuracy of CAS is also comparable to other psychiatric screening tools (Evren et al., 2020).

The results demonstrated that the CVAS Teen Version had significant positive correlation with all five sub-dimensions, which indicates that the scale can detect whether respondents indeed have vaccination anxiety as well as significant emotional, cognitive, physiological, behavioral, and emotional regulation symptoms, which have more deleterious effects on the psychological health of teens. These findings are also in tune with other studies, such as Ahorsu et al. (2020), Harper et al. (2020), and Lee (2020).

Both sub-dimensions of Emotion Regulation had significant positive associations with other dimensions of the CVAS Teen Version. Difficulties

in emotional regulation are important to detect by health care professionals. Gross and Jazaieri (2014) reported that difficulty in regulating emotion is the inability to control emotional responses to the stimuli that produce those emotions and such individuals experience emotional states in maladaptive ways. Silk et al. (2003) stated that emotion regulation problems have been identified in more than half of the first and second axis disorders, which was also shown in the research study of Repetti et al. (2002). According to Ochsner and Gross (2007), many psychiatric disorders are associated with emotional instability and emotional dysregulation. Emotional dysregulation is particularly strongly correlated with depression and anxiety disorders (Phillips et al., 2002; Mennin, 2004).

Higher CVAS Teen Version scores are associated with negative psychological effects and maladaptive strategies, such as difficulty regulating emotion when experiencing coronavirus vaccination anxiety (Mohammadpour et al., 2020). As already discussed, many suffer anxiety about the COVID-19 disease; for example, health care workers have been reported to experience anxiety due to exposure to patients suspected or confirmed to have COVID-19 (Lai et al., 2020). Nonetheless, such individuals may also be afraid and anxious concerning the effectiveness of COVID-19 vaccinations (Lee, 2020a). It can be assumed that people who cannot control their emotions regarding COVID-19 may experience anxiety regarding vaccinations. And those individuals have a strong impact on their younger family members, including teens and adolescents, leading them to decline vaccines or develop more fear and anxiety regarding the consequences of COVID-19 vaccinations.

The divergent validity of the scale was also evaluated. According to the results, the highest HTMT score was 0.694, between the Behavioral

Dimension and Suppression, one of the sub-domains of Emotion Regulation, and because this was less than the maximum tolerable threshold of 0.85, this confirms that the discriminant validity was not violated. Similarly, the CFA model fit met the criteria and did not violate the threshold, which served as validation of this model. These findings indicate that individuals in the age range of 12-18 are less flexible and will have more anxiety regarding Covid vaccination. While in such a situation, the age group of 12-18 is more vulnerable to develop symptoms and psychological issues due to fear and anxiety regarding the vaccinations' effects (Barth, 2021). Studies have shown that individuals with high avoidance experience more emotional responses to stimuli and report severe symptoms of stress and anxiety (Tull et al., 2004; Solan, 2004). These findings are consistent with the outcomes of other research studies that have examined the psychological effects of the COVID-19 pandemic, including Zhang et al. (2020), Liu et al. (2020), and Li et al. (2020).

### **Limitations and Suggestions**

There are also some limitations to this study. First, in the current research, participants belonged to the general Egyptian teen population and no formal diagnoses regarding mood or mental disorders were considered (e.g., generalized anxiety, stress, etc.). Second, there is no doubt that there is subjectivity involved in anxiety and it not easy to assess objectivity. Third, we used a convenient sampling method, which may affect the generalizability of the results.

It is suggested that in future studies, this tool be tested in individuals known to have hesitancy and fear and anxieties regarding COVID-19 vaccinations in a large sample to determine its sensitivity and specificity. A

further suggestion is that to overcome fake news about the vaccines, which trigger anxieties, it is important to provide reliable and credible sources of information, and public awareness programs should be developed to disseminate the facts regarding the safety of the vaccines.

## **Conclusion**

The present study confirmed that the CVAS Teen Version, based on five dimensions, has good psychometric properties that can be used to assess anxieties among teens regarding COVID-19 vaccination. Therefore, therapists, health professionals, and researchers can use this screening tool to diagnose COVID-19 vaccination anxiety in individuals aged between 12 and 18.

## **Acknowledgments**

This project was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah (*Grant Z: 9-246-1440*). The authors, therefore, acknowledge with thanks DSR technical and financial support.

## REFERENCES

- Ahorsu, D. K., Lin, C. Y., Imani, V., Saffari, M., Griffiths, M. D., & Pakpour, A. H. (2020). The Fear of COVID-19 Scale: Development and initial validation. *International Journal of Mental Health and Addiction*, 1-9. Advance online publication. <https://doi.org/10.1007/s11469-020-00270-8>
- Ali, A. M., & Hassan, K. A. (2023). The role of social media in raising awareness about the COVID-19 pandemic. *Sharjah Journal of Humanities*, 30(2), 123-145. <https://spu.sharjah.ac.ae/index.php/HSS/article/view/1761>
- Ali, N. (2024). The role of evidence-based practice in clinical social work in developing the professional performance of workers with COVID-19 patients. *University of Sharjah Journal of Humanities and Social Sciences*, 15(4), 525-535. Retrieved from <https://spu.sharjah.ac.ae/index.php/HSS/article/view/4103>
- Arafa, A., Mohammed, Z., Mahmoud, O., Elshazley, M., & Ewis, A. (2021). Depressed, anxious, and stressed: What have healthcare workers on the frontlines in Egypt and Saudi Arabia experienced during the COVID-19 pandemic? *Journal of Affective Disorders*, 278, 365-371.
- Arbuckle, J. L. (2016). *IBM SPSS Amos 24 user's guide*. IBM.
- Asmundson, G. J., & Taylor, S. (2020). Coronaphobia: Fear and the 2019-nCoV outbreak. *Journal of Anxiety Disorders*, 70, 102196. <https://doi.org/10.1016/j.janxdis.2020.102196>
- Atiyat, A., & Abu Hamour, A. (2022). Evaluation of distance education services for students of special education centers in Jordan during the COVID-19 pandemic. *University of Sharjah Journal of Humanities and Social Sciences*, 19(1), 497-530. <https://doi.org/10.36394/jhss/19/1/19>
- Bajema, K. L., Oster, A. M., McGovern, O. L., Lindstrom, S., Stenger, M. R., Anderson, T. C., ... & 2019-nCoV Persons Under Investigation Team (2020). Persons evaluated for 2019 Novel Coronavirus - United States, January 2020. *MMWR Morbidity and Mortality Weekly Report*, 69(6), 166-170. <https://doi.org/10.15585/mmwr.mm6906e1>
- Bandalos, D. L., & Finney, S. J. (2010). Factor analysis: Exploratory and confirmatory. In G. R. Hancock & R. O. Mueller (Eds.), *The reviewer's guide to quantitative methods in the social sciences* (pp. 93-114). New York: Routledge.
- Banerjee, D. (2020). The COVID-19 outbreak: Crucial role the psychiatrists can play. *Asian Journal of Psychiatry*, 50, 102014.
- Barth, E. (2021, May 26). COVID-19 vaccine anxiety among teens. *Allina Health*. <https://www.allinahealth.org/allina-news/2021/05/covid19-vaccine-anxiety-among-teens>
- Bodner, E., Bergman, Y. S., Ben-David, B., & Palgi, Y. (2021). Vaccination anxiety when vaccinations are available: The role of existential concerns. *Stress and Health*, 1-8. <https://doi.org/10.1002/smi.3079>

- Boomsma, A., Hoyle, R. H., & Panter, A. T. (2012). The structural equation modeling research report. In R. H. Hoyle (Ed.), *Handbook of structural equation modeling* (pp. 341–358). New York: Guilford Press.
- Brenner, H., & Kliebsch, U. (1996). Dependence of weighted kappa coefficients on the number of categories. *Epidemiology*, 7(2), 199-202.
- Brown, T. A. (2015). *Confirmatory Factor Analysis for applied research (2nd ed.)*. New York: Guilford Publications.
- Byrne, B. M. (2016). *Structural equation modeling with AMOS: Basic concepts, applications, and programming (3rd ed.)*. Mahwah, NJ: Erlbaum.
- Chan, J. F. W., Yip, C. C. Y., To, K. K. W., Tang, T. H. C., Wong, S. C. Y., Leung, K. H., ... & Yuen, K. Y. (2020). Improved molecular diagnosis of COVID-19 by the novel, highly sensitive and specific COVID-19-RdRp/Hel real-time reverse transcription-PCR assay validated in vitro and with clinical specimens. *Journal of Clinical Microbiology*, 58(5), e00310-20.
- Chong, M. Y., Wang, W. C., Hsieh, W. C., Lee, C. Y., Chiu, N. M., Yeh, W. C., ... & Chen, C. L. (2004). Psychological impact of severe acute respiratory syndrome on health workers in a tertiary hospital. *The British Journal of Psychiatry*, 185(2), 127-133.
- Daly, M., & Robinson, E. (2020). Willingness to vaccinate against COVID-19 in the US: Longitudinal evidence from a nationally representative sample of adults from April-October 2020. *medRxiv*. <https://doi.org/10.1101/2020.11.27.20239970>
- Detoc, M., Bruel, S., Frappe, P., Tardy, B., Botelho-Nevers, E., & Gagneux-Brunon, A. (2020). Intention to participate in a COVID-19 vaccine clinical trial and to get vaccinated against COVID-19 in France during the pandemic. *Vaccine*, 38(45), 7002-7006.
- Dong, X. Y., Wang, L., Tao, Y. X., Suo, X. L., Li, Y. C., Liu, F., ... & Zhang, Q. (2017). Psychometric properties of the Anxiety Inventory for Respiratory Disease in patients with COPD in China. *International Journal of Chronic Obstructive Pulmonary Disease*, 12, 49.
- Esteban, R. F. C., Mejia, C. R., Rodriguez-Alarcon, J. F., Ticona, D., Flores-Lovon, K., Paredes-Obando, M., ... & Tovani-Palone, M. R. (2021). Validation of a scale to measure the perception of SARS-CoV-2 vaccines acceptance: The VAC-COVID-19 Scalescale. *Electronic Journal of General Medicine*, 18(5), em303. <https://doi.org/10.29333/ejgm/11012>
- Evren, C., Evren, B., Dalbudak, E., Topçu, M., & Kutlu, N. (2020). Measuring anxiety related to COVID-19: A Turkish validation study of the Coronavirus Anxiety Scale. *Death Studies*, June 3, 1-7. <https://doi.org/10.1080/07481187.2020.1774969>
- Fox, J., Byrnes, J. E., Boker, S., & Neale, M. C. (2012). Structural equation modeling in R with the sem and OpenMx packages. In R. H. Hoyle (Ed.), *Handbook of Structural Equation Modeling* (pp. 325–340). New York: Guilford Press.

- Gana, K., & Broc, G. (2019). *Structural equation modeling with lavaan*. London: Wiley.
- Gefen, D., Rigdon, E. E., & Straub, D. (2011). An update and extension to SEM guidelines for administrative and social science research. *Management Information Systems Quarterly*, 35, A1-A7.
- Hamza Shuja, K., Aqeel, M., Jaffar, A., & Ahmed, A. (2020). COVID-19 pandemic and impending global mental health implications. *Psychiatria Danubina*, 32(1), 32-35.
- Hancock, G. R., & Mueller, R. O. (Eds.). (2013). *Structural equation modeling: A second course (2nd ed.)*. Charlotte, NC: Information Age Publishing, Inc.
- Harper, C. A., Satchell, L. P., Fido, D., & Latzman, R. D. (2020). Functional fear predicts public health compliance in the COVID-19 pandemic. *International Journal of Mental Health and Addiction*, 1-14. <https://doi.org/10.1007/s11469-020-00281-5>
- Hause, A. M., Gee, J., Johnson, T., Jazwa, A., Marquez, P., Miller, E., ... & Shay, D. K. (2021). Anxiety-related adverse event clusters after Janssen COVID-19 vaccination—five U.S. mass vaccination sites, April 2021. *MMWR Morb Mortal Wkly Rep* 2021, 70, 685–688. <https://doi.org/10.15585/mmwr.mm7018e3>
- Howitt, D., & Cramer, D. (2017). *Understanding statistics in psychology with SPSS*. Boston: Pearson Higher Education.
- Hox, J. J. (2013). Multilevel regression and multilevel structural equation modeling. In T. D. Little (Ed.), *The Oxford Handbook of Quantitative Methods* (pp. 281-294). New York: Oxford University Press.
- Hoyle, R. H. (Ed.). (2012). *Handbook of structural equation modeling*. New York: Guilford Press.
- Judd, C. M., McClelland, G. H., & Ryan, C. S. (2017). *Data analysis: A model comparison approach*. Routledge: New York.
- Kilic, A. (2018). Can factor scores be used instead of total score and ability estimation? *International Journal of Assessment Tools in Education*, 6(1), 25-35.
- Kline, R. B. (2016). *Principles and practice of structural equation modeling (4th edition)*. New York/London: Guilford.
- Khusaifan, S. J., & Samak, Y. A. (2024). Psychometric properties of the COVID-19 vaccine anxiety scale for adults. *Journal of Environmental and Agricultural Health Sciences*, 37(1), 524. <https://doi.org/10.33193/JEAHS.37.2024.524>
- Kumar, A., & Somani, A. (2020). Dealing with Corona virus anxiety and OCD. *Asian Journal of Psychiatry*, 51, 102053. <https://doi.org/10.1016/j.ajp.2020.102053>
- Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., ... & Hu, S. (2020). Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Network Open*, 3(3), e203976-e203976. <https://doi.org/10.1001/jamanetworkopen.2020.3976>

- Lazarus, J. V., Ratzan, S. C., Palayew, A., Gostin, L. O., Larson, H. J., Rabin, K., ... & El-Mohandes, A. (2021). A global survey of potential acceptance of a COVID-19 vaccine. *Nature Medicine*, 27(2), 225-228.
- Lee, S. A. (2020a). Coronavirus Anxiety Scale: A brief mental health screener for COVID-19 related anxiety. *Death Studies*, 44(7), 393-401.
- Lee, S. A. (2020b). How much "Thinking" about COVID-19 is clinically dysfunctional? *Brain, Behavior, and Immunity*, 87, 97.
- Li, S., Wang, Y., Xue, J., Zhao, N., & Zhu, T. (2020). The impact of COVID-19 epidemic declaration on psychological consequences: A study on active Weibo users. *International Journal of Environmental Research and Public Health*, 17(6), 2032. <https://doi.org/10.3390/ijerph17062032>
- Little, J. (2013). Multilevel confirmatory ordinal factor analysis of the Life Skills Profile-16. *Psychological Assessment*, 25, 810-825.
- Liu, N., Zhang, F., Wei, C., Jia, Y., Shang, Z., Sun, L., ... & Liu, W. (2020). Prevalence and predictors of PTSS during COVID-19 outbreak in China hardest-hit areas: Gender differences matter. *Psychiatry Research*, 287, 112921. <https://doi.org/10.1016/j.psychres.2020.112921>
- Loehlin, J. C., & Beaujean, A. A. (2017). *Latent variable models: An introduction to factor, path, and structural equation analysis*. New York, NY: Taylor & Francis.
- Lurie, N., Saville, M., Hatchett, R., & Halton, J. (2020). Developing COVID-19 vaccines at pandemic speed. *New England Journal of Medicine*, 382(21), 1969-1973.
- Magadmi, R. M., & Kamel, F. O. (2021). Beliefs and barriers associated with COVID-19 vaccination among the general population in Saudi Arabia. *BMC Public Health*, 21(1), 1-8.
- Malik, A., Malik, J., & Ishaq, U. (2021). Acceptance of COVID-19 vaccine in Pakistan among health care workers. *medRxiv*. <https://doi.org/10.1101/2021.02.23.21252271>
- Mohammadpour, M., Ghorbani, V., Khoramnia, S., Ahmadi, S. M., Ghvami, M., & Maleki, M. (2020). Anxiety, self-compassion, gender differences and COVID-19: Predicting self-care behaviors and fear of COVID-19 based on anxiety and self-compassion with an emphasis on gender differences. *Iranian Journal of Psychiatry*, 15(3), 213. <https://doi.org/10.18502/ijps.v15i3.3813>
- Murray, C. J., & Piot, P. (2021). The potential future of the COVID-19 pandemic: will SARS-CoV-2 become a recurrent seasonal infection? *JAMA*, 325(13), 1249-1250.
- Neumann-Böhme, S., Varghese, N. E., Sabat, I., Barros, P. P., Brouwer, W., van Exel, J., ... & Stargardt, T. (2020). Once we have it, will we use it? A European survey on willingness to be vaccinated against COVID-19. *The European Journal of Health Economics*, 21(7), 977-982. <https://doi.org/10.1007/s10198-020-01208-6>

- Nzaji, M. K., Ngombe, L. K., Mwamba, G. N., Ndala, D. B. B., Miema, J. M., Lungoyo, C. L., ... & Musenga, E. M. (2020). Acceptability of vaccination against COVID-19 among healthcare workers in the Democratic Republic of the Congo. *Pragmatic and Observational Research, 11*, 103.
- Pérez-Fuentes, M. D. C., Molero Jurado, M. D. M., Oropesa Ruiz, N. F., Martos Martínez, Á., Simón Márquez, M. D. M., Herrera-Peco, I., & Gázquez Linares, J. J. (2020). Questionnaire on perception of threat from COVID-19. *Journal of Clinical Medicine, 9*(4), 1196.
- Pogue, K., Jensen, J. L., Stancil, C. K., Ferguson, D. G., Hughes, S. J., Mello, E. J., ... & Poole, B. D. (2020). Influences on attitudes regarding potential COVID-19 vaccination in the United States. *Vaccines, 8*(4), 582.
- Portney, L. G., & Watkins, M. P. (2009). *Foundations of clinical research: Applications to practice* (Vol. 892). Upper Saddle River, NJ: Pearson/Prentice Hall.
- Preacher, K. J., & MacCallum, R. C. (2002). Exploratory factor analysis in behavior genetics research: Factor recovery with small sample sizes. *Behavior Genetics, 32*(2), 153-161.
- Saied, S. M., Saied, E. M., Kabbash, I. A., & Abdo, S. A. E. F. (2021). Vaccine hesitancy: Beliefs and barriers associated with COVID-19 vaccination among Egyptian medical students. *Journal of Medical Virology, 93*(7), 4280-4291.
- Sallam, M., Dababseh, D., Eid, H., Al-Mahzoum, K., Al-Haidar, A., Taim, D., ... & Mahafzah, A. (2021). High rates of COVID-19 vaccine hesitancy and its association with conspiracy beliefs: A study in Jordan and Kuwait among other Arab countries. *Vaccines, 9*(1), 42.
- Serrano-Mollar, A. (2012). Alveolar epithelial cell injury as an etiopathogenic factor in pulmonary fibrosis. *Archivos de bronconeumologia, 48*, 2-6.
- Sloan, D. M. (2004). Emotion regulation in action: Emotional reactivity in experiential avoidance. *Behaviour Research and Therapy, 42*(11), 1257-1270.
- Szilagyi, P. G., Thomas, K., Shah, M. D., Vizueta, N., Cui, Y., Vangala, S., & Kapteyn, A. (2021). National trends in the US public's likelihood of getting a COVID-19 vaccine—April 1 to December 8, 2020. *JAMA, 325*(4), 396-398.
- Taylor, S. (2019). *The psychology of pandemics: Preparing for the next global outbreak of infectious disease*. Cambridge Scholars Publishing.
- Thompson, B. (2018). *Exploratory and confirmatory factor analysis: Understanding concepts and applications*. Washington, DC: American Psychological Association.
- Tull, M. T., Gratz, K. L., Salters, K., & Roemer, L. (2004). The role of experiential avoidance in posttraumatic stress symptoms and symptoms of depression, anxiety, and somatization. *The Journal of Nervous and Mental Disease, 192*(11), 754-761.

- Wang, J., & Wang, X. (2019). *Structural equation modeling: Applications using Mplus*. Chichester, West Sussex: John Wiley.
- Wang, J., Jing, R., Lai, X., Zhang, H., Lyu, Y., Knoll, M. D., & Fang, H. (2020). Acceptance of COVID-19 vaccination during the COVID-19 pandemic in China. *Vaccines*, 8(3), 482.
- Wheaton, M. G., Abramowitz, J. S., Berman, N. C., Fabricant, L. E., & Olatunji, B. O. (2012). Psychological predictors of anxiety in response to the H1N1 (swine flu) pandemic. *Cognitive Therapy and Research*, 36(3), 210-218.
- WHO (World Health Organization) (2021). WHO Director-General's opening remarks at the media briefing on COVID-19-4 August 2021. Retrieved from <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-4-august-2021>.
- Wu, P., Fang, Y., Guan, Z., Fan, B., Kong, J., Yao, Z., ... & Hoven, C. W. (2009). The psychological impact of the SARS epidemic on hospital employees in China: Exposure, risk perception, and altruistic acceptance of risk. *The Canadian Journal of Psychiatry*, 54(5), 302-311.
- Yip, P. S., Cheung, Y. T., Chau, P. H., & Law, Y. W. (2010). The impact of epidemic outbreak. *Crisis*, 31(2). <https://doi.org/10.1027/0227-5910/a000015>
- Yohannes, A. M., Willgoss, T. G., Fatoye, F. A., & Goldbart, J. (2013). Validity and reliability of the anxiety inventory for respiratory disease scale in patients with COPD. *American Thoracic Society*, 187(1). [https://doi.org/10.1164/ajrccm-conference.2013.187.1\\_MeetingAbstracts.A4227](https://doi.org/10.1164/ajrccm-conference.2013.187.1_MeetingAbstracts.A4227).
- Zhang, C., Yang, L., Liu, S., Ma, S., Wang, Y., Cai, Z., ... & Zhang, B. (2020). Survey of insomnia and related social psychological factors among medical staff involved in the 2019 novel coronavirus disease outbreak. *Frontiers in Psychiatry*, 11, 306.

## Appendix

### Final Version of Covid Vaccination Anxiety Teen Version

	Strongly Disagree	Disagree	Neutral	Agree	Strongly Agree
1. I find it difficult to relax when I think about taking the COVID-19 vaccine.					
2. The thought of getting a vaccination makes me restless.					
3. Even after information about vaccine trials and success rates, I am not able to control my worry.					
4. I am afraid that the vaccine could cause possible harm to me and my family.					
5. I am scared that the COVID-19 vaccine could cause the disease.					
6. I use to have repetitive negative thoughts about the consequences of COVID					
7. I am fearful of the side effects of the COVID-19 vaccine.					
8. I am scared that the COVID-19 vaccine could cause some sort of deformity or dysfunction in my body.					
9. I experience shortness of breath when I think of taking COVID					

<b>10.</b> I experience chest discomfort when I read or listen to COVID-19 vaccine news.					
<b>11.</b> I feel nauseous when I think of getting the COVID-19 vaccine.					
<b>12.</b> I experience loosening of bowels when think of taking the COVID-19 vaccine.					
<b>13.</b> My heart starts to run faster when I think or read about the COVID-19 vaccine.					
<b>14.</b> I feel some sensation in my muscles when I think about getting the COVID-19 vaccine.					
<b>15.</b> I experience pain in my muscles when I think of getting the COVID-19 vaccine.					
<b>16.</b> I experience hot and cold flushes when I think about getting the COVID-19 vaccine.					
<b>17.</b> I feel numb when I think of taking the COVID-19 vaccine.					
<b>18.</b> I frequently check for vaccine validation facts to ensure its effectiveness.					
<b>19.</b> The thought of taking the COVID 19 vaccine causes difficulty in falling asleep, broken sleep, and 2 bad dreams.					

<b>20.</b> I would prefer to avoid vaccines in case it's not compulsory					
<b>21.</b> When I want to feel happier I think about something different					
<b>22.</b> When I want to feel less bad (e.g., sad, angry, worried) I think about something different.					
<b>23.</b> When I'm worried about something; I make myself think about it in a way that helps me feel better.					
<b>24.</b> When I want to feel happier about something, I change the way I'm thinking about it.					
<b>25.</b> I control my feelings about things by changing the way I think about them.					
<b>26.</b> When I want to feel less bad (e.g., sad, angry, worried )about something I change the way I'm thinking about it.					
<b>27.</b> I keep my feelings to myself.					
<b>28.</b> When I am feeling happy, I am careful not to show it.					
<b>29.</b> I control my feelings by not showing them.					
<b>30.</b> When I'm feeling bad (e.g., sad, angry, worried ), I am careful not to show it.					

### **List of Abbreviations in the Manuscript:**

**COVID-19** - Coronavirus Disease 2019

**CFA** - Confirmatory Factor Analysis

**CMIN** - Minimum Discrepancy (Chi-square fit statistic)

**DF** - Degrees of Freedom

**P** - Probability Value

**CMIN/DF** - Chi-square divided by Degrees of Freedom (Model Fit Index)

**NFI** - Normed Fit Index

**RFI** - Relative Fit Index

**IFI** - Incremental Fit Index

**TLI** - Tucker-Lewis Index

**CFI** - Comparative Fit Index

**PRATIO** - Parsimony Ratio

**PNFI** - Parsimony Normed Fit Index

**PCFI** - Parsimony Comparative Fit Index

**RMSEA** - Root Mean Square Error of Approximation

**LO 90** - Lower Bound of 90% Confidence Interval for RMSEA

**HI 90** - Upper Bound of 90% Confidence Interval for RMSEA

**PCLOSE** - Probability of Close Fit

**EFA** - Exploratory Factor Analysis

**KMO** - Kaiser-Meyer-Olkin Measure of Sampling Adequacy

**ICC** - Intraclass Correlation Coefficient

**KC** - Kappa Coefficient

**AVE** - Average Variance Extracted

**CR** - Composite Reliability

**MSV** - Maximum Shared Variance

**HTMT** - Heterotrait-Monotrait Ratio

**PD** - Physiological Dimension

**CD** - Cognitive Dimension

**ED** - Emotional Dimension

**BD** - Behavioral Dimension

**ER** - Emotion Regulation

**RE** - Reappraisal

**SUP** - Suppression

**WHO** - World Health Organization

**PTSD** - Post-Traumatic Stress Disorder

**SPSS** - Statistical Package for the Social Sciences

## مقياس قلق لقاح كوفيد - 19 للمراهقين: مقياس لتقييم القلق المرتبط بلقاحات كوفيد - 19 بين المراهقين في مصر

نسرين يعقوب محمد خليل<sup>(1)</sup>

ياسر عبد العظيم عبد الموجود سمك<sup>(2)</sup>

### ملخص البحث:

الهدف: كان لجائحة كوفيد-19- تأثيرات هائلة على الحياة الجسدية والصحة النفسية للأشخاص في جميع أنحاء العالم. لم يتأثر البالغون فقط، بل عانى الأطفال والمراهقون أيضًا. وقد أثير الأمل بالعودة إلى الحياة الطبيعية مع تطوير اللقاحات، لكن انتشار المعلومات المضللة بشأن فعالية وسلامة اللقاحات خلق ظاهرة نفسية جديدة، وهي القلق بشأن اللقاحات نفسها. الأطفال أكثر عرضة للتأثر بمثل هذه المخاوف والقلق. لمعالجة هذه المشكلة، تم تطوير مقياس فريد لقياس القلق بين المراهقين بشأن لقاحات كوفيد-19-، وتم تقييم الخصائص السيكومترية للمقياس

المنهجية: تم تطوير عناصر المقياس وبعد الوصول إلى 30 عنصرًا إجماليًا، تم تجميع العناصر وتقسيمها إلى خمسة أبعاد فرعية. تقيس هذه الأبعاد الجوانب: العاطفية، والمعرفية، والفسولوجية، والسلوكية، وتنظيم الانفعالات. تم جمع البيانات من 1296 مشاركًا تتراوح أعمارهم بين 12-18 عامًا. للتحقق من الخصائص السيكومترية للمقياس، تم إجراء تحليل العوامل الاستكشافية، وتحليل التباين، ومصنوفة العوامل المدورة، والصلاحية الإنشائية والموثوقية، بالإضافة إلى تحليل العوامل التأكيدية

النتائج: بالنسبة للأبعاد الخمسة التي تم تطويرها، كانت معاملات ألفا كرونباخ لموثوقية المقياس كالتالي: البعد العاطفي، 0.797؛ والبعد المعرفي، 0.857؛ والبعد الفسولوجي، 0.951؛ والبعد السلوكي، 0.769؛ وتنظيم الانفعالات، 0.900.

الخلاصة: المقياس المطور موثوق وصالح لقياس قلق اللقاحات بين المراهقين. يجب أن تتابع الدراسات المستقبلية هذه الدراسة مع عينة أكبر من السكان على مستوى العالم

الكلمات الدالة: مقياس قلق لقاح كوفيد-19-، نسخة المراهقين لمقياس قلق لقاح كوفيد-19-، اللقاح، القلق، الصدق، الصلاحية، مصر

(1) كلية الآداب والعلوم الإنسانية - جامعة الملك عبد العزيز (جدة - المملكة العربية السعودية) geopsychologie@gmail.com

(2) كلية الآداب - جامعة أسيوط (أسيوط - جمهورية مصر العربية)